



The deep blue day is decreasing in China

Su Wang^{1,2} · Gang Huang^{1,2,3} · Kaiming Hu^{1,4} · Lin Wang⁵ · Tie Dai^{1,6} · Chunjiang Zhou^{1,2}

Received: 27 June 2021 / Accepted: 11 December 2021 / Published online: 22 January 2022

© The Author(s), under exclusive licence to Springer-Verlag GmbH Austria, part of Springer Nature 2021

Abstract

The deep blue sky is an indicator of a lower concentration of aerosols and a cloudless sky. With increasing human emissions, a trend towards days with fewer deep blue skies might indicate a decline in a good living environment for humans. This study investigates the long-term changes of the deep blue sky in China from 1980 to 2018. Due to a lack of direct measurements, we use atmospheric visibility and low cloud cover to classify blue sky days into three grades: light blue day, medium blue day, and deep blue day. Climatologically, annual deep blue days increase from southeast China to northwest China, with the maximum number in Xinjiang and eastern Inner Mongolia and the minimum number in western Qinghai and southern Hebei. From 1980 to 2018, annual deep blue days show a prominent decreasing trend in most of China, with area-mean annual deep blue days decreasing by -0.48 days per year (d/y) in China, and the variation becomes more obvious after 2013. The maximum decreasing trend is observed in eastern China. The most prominent decreases of deep blue days are seen in winter. Both air pollution and the change in meteorological conditions contribute to the decrease of wintertime deep blue days in China. Specifically, the decrease in surface wind speed hinders the cleaning of air by winds, the increase in surface air temperature, and decrease in relative humidity is favorable for low cloud increase, and the increasing emission of pollution reduces atmospheric visibility.

1 Introduction

With the improvement in living standards, there is a growing desire for a better atmospheric environment and more clear blue skies. Traditionally, a sunny day with few clouds means blue sky day. However, with increasing anthropogenic emissions, the sky might be less or not blue even when the weather condition is “sunny (defined by cloud cover)” (Fast et al. 2006; Yang et al. 2011; Lin et al. 2012; Li et al. 2016a). On the other hand, fewer air pollutions do not directly bring in more blue skies, considering days with higher cloud cover or precipitation. The inconsistency between dropping PM and unchanged low visibility events (Ding et al. 2016) can be seen even during the COVID-19 lockdown period (Huang et al. 2020b; Huang et al. 2020c; Liu et al. 2020). All mentioned above try to convey the non-synergy correlation between air pollution and blue days when considering the unfavorable meteorological conditions dominated by large-scale synoptic circulation (Bei et al. 2020a). Thus, there are reasons to believe that “sunny” or “clean” may not be sufficient for a blue sky.

A series of heavy air-pollution events in the early 2010s in China have raised people’s strong desire for better air quality as well as more blue skies (Zhang et al. 2014; Zhang

✉ Gang Huang
hg@mail.iap.ac.cn

✉ Kaiming Hu
hkm@mail.iap.ac.cn

¹ State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

² University of Chinese Academy of Sciences, Beijing 100049, China

³ Laboratory for Regional Oceanography and Numerical Modeling, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China

⁴ Center for Monsoon System Research, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

⁵ Key Laboratory of Regional Climate-Environment for Temperate East Asia, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

⁶ Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters/Key Laboratory of Meteorological Disaster of Ministry of Education, Nanjing University of Information Science and Technology, Nanjing, China

et al. 2015). The State Council of China has issued a series of strict anti-pollution control, such as the Air Pollution Prevention and Control Action Plan (China State Council 2013) and Blue-Sky Protection Campaign (China State Council 2018). However, there may be a discrepancy about whether clean actions have brought in more blue days by studies focusing on air quality or cloud cover (Zhang et al. 2019c; Bei et al. 2020b; Dai et al. 2020). Therefore, lacking direct index, it is still a challenge to understand the change of blue sky days in China and its underlying mechanism.

The first challenge is how to define a blue sky day. The occurrence of the blue sky depends not only on meteorological conditions but also on environmental conditions. First, a blue sky day should be a sunny day, which can be identified by low cloud cover and precipitation records. Second, a blue sky day should have clean air. Previous studies show that good air quality can be represented by high atmospheric visibility in the long term (Wu 2004; Ma et al. 2014; Zhang et al. 2014; Liu et al. 2017). Thus, our previous study (Wang et al. 2019) defined a blue sky day via the following criteria: the days with no rain, low cloud cover ≤ 75 th percentile, and the dry visibility in 14:00 ≥ 15 km.

However, the above definition does not tell the degree of blue of a blue sky day. In the real world, the gradations of the blue sky days are different from each other via presenting different RGB values (Fig. S1), which is dominated by the discrepancy between meteorology and the environment. A continuous deep blue days during the APEC (Asia-Pacific Economic Cooperation) conference in Beijing during December 2014 (Liu et al. 2016; Sun et al. 2016; Zhang et al. 2016; Xu et al. 2018) attract people's attention around the world. Thus, classifying the blue days into different degrees will help us understand the change of blue days better.

The second challenge is how to assess the relative contributions of meteorological conditions and environmental factors to the change in the blue sky days. Observed evidence (Panmao et al. 1999; Alexander et al. 2006; Zhou et al. 2009; Hu et al. 2013; Easterling et al. 2019) shows that East Asia has experienced prominent changes in rainfall, temperature, and atmospheric circulation during recent decades. Meanwhile, with the development of industry, the emissions of aerosol have also increased significantly (Kato and Akimoto 2007; Meinshausen et al. 2009; Zhang and Wei 2014; Millar et al. 2017). Many studies show that the changes in meteorological and environmental conditions are responsible for the increase of air pollution events in China during recent decades (Kaiser 2002; Li et al. 2014; Dufour et al. 2015; Dai et al. 2020). For example, the weakened monsoon may increase the aerosol concentrations over eastern China (Zhu et al. 2012; Li et al. 2016b) and amplify persistent haze in Beijing (Li et al. 2016b; Cai et al. 2017). Nevertheless, we still do not know the role of meteorological and environmental factors in blue sky day change in China.

This study aims to creatively classify blue days into 3 grades, analyze their spatial-temporal variations during 1980–2018 in China, and reveal the effect of air pollution prevention plans. The rest of the paper is organized as follows. Section 2 describes the data and methods used. Section 3 shows evidence for deep blue sky days decrease in China especially in winter and discusses the possible reasons for the change. Section 4 includes a conclusion and some discussions.

The classified blue days could be a direct index to evaluate the efficiency of the pollution prevention and control measures quantitatively in recent years. As life satisfaction and happiness are closely related to deep blue sky (Welsch 2006; Luechinger 2010; Li et al. 2014), this work will help to the construction of people's livelihood. Meanwhile, it could provide a reference to studies on the relationship between weather and disease such as lung cancer and abortion (Pope et al. 2002; Zhang et al. 2019a; Zhang et al. 2019b). Focusing on days suitable for outdoor activities, it also serves the tourist industry and selection of important events. Finally, studying the mechanism behind the deep blue days will be a referee to host mega-events like Olympic Winter Games and implement policies on energy conservation and emission reduction.

2 Data and methods

2.1 Data

The daily meteorological observation data, including wind speed at 10 m, relative humidity, low cloud cover, and visibility values at 14:00 BJT from 1980 to 2018, were derived from the National Meteorological Information Center of the China Meteorological Administration. The monthly European Centre for Medium-Range Weather Forecasts (ECMWF) Re-Analysis-Interim atmospheric reanalysis data from January 1980 to December 2018 with a horizontal resolution of $0.75^\circ \times 0.75^\circ$ (Dee et al. 2011) are used in the analysis of climatic conditions for blue sky days. The Community Emissions Data System (CEDS) for historical emissions from 1980 to 2014 (Hoesly et al. 2018) is used as an auxiliary to study the anthropogenic contribution, which is available from <https://github.com/JGCRI/CEDS/>.

2.2 Data quality controls

To ensure the accuracy and continuity of data used, a series of strict quality controls have been implemented following earlier work (Wang et al. 2019). In particular, as manual visibility measurement is broadly replaced by automated detection techniques around 2013–2015, special treatment is required for visibility records to keep data consistency. To this end, we transform the visibility during the post-2013 period according to the formula below

(Organization 2008; Pei et al. 2018; Zhang et al. 2020a) to maintain data consistency:

$$\frac{VIS_{auto}}{VIS_{manual}} = \frac{(1/k) \times \ln(1/0.05)}{(1/k) \times \ln(1/0.02)} \approx 0.766$$

where k represents the extinction coefficient, and 0.05 and 0.02 denote the thresholds taken by the automatic and human observer, respectively. The $visibility_{manual}$ and $visibility_{auto}$ are the human and instrument measurements of atmospheric visibility, respectively. In the analyses, the maximum of human observing visibility has been set to 30 km to reduce the discrepancy between the two observed methods. We also use the Standard Normal Homogeneity Test (SNHT) (Alexandersson and Moberg 1997) to verify whether stations exhibit considerable discontinuity in annual average visibility, by which only station observations passing the 99% confidence level test have been retained. Earlier studies have compared manual and automatic observations (Fan et al. 2017; Chang et al. 2020; Zhang et al. 2020a), and their results reported that although the threshold has been modified during 2013–2015, the continuity of the long-term data is not affected (Yin et al. 2017; Pei et al. 2018; Wang et al. 2020). Recent work demonstrated that this data adjustment can improve the continuity and reliability of visibility data (Zhang et al. 2020a). After quality control, data at 378 monitoring stations remain for further investigation (Fig. 1).

3 Methods

3.1 Classification of blue days

The classification procedures of blue days are as follows. Following our earlier work that takes the low cloud cover of the individual station into consideration, two thresholds are selected to represent the blue days at different levels. One is the national mean visibility (21.52 km) in all blue

days from 1980 to 2018 in China (T1), and the other is the multi-year average value of low cloud cover in blue days at each station (T2). With the defined thresholds, we build two corresponding criteria to check: (1) whether the visibility is larger than T1; (2) whether the low cloud cover is less than T2. Based upon the above two criteria, Chinese blue days (CBDs) can be divided into three categories.

- (1) Deep blue: If one day meets both two prerequisites, it is recognized as the deep blue day (CBD3), representing the most desirable rate of blue days for people.
- (2) Medium blue: If it fits only one of them, it is termed as medium blue days (CBD2).
- (3) Light blue (CBD1): If it satisfies neither of the two prerequisites, it is ranked as the light blue days (CBD1), suggesting that it merely reaches the lowest level: the weakest blue days.

Case studies have been conducted to test the accuracy of the classification. Almost all well-known blue events, such as the APEC blue and Parade blue fall into the CBD3 category (not shown), indicating that our classification is reasonable.

4 Methods

A linear-trend analysis is used to reveal the variations of the time series of variables. Pearson correlation, and the standardized multivariate linear regression model (SMLR), and the Student's t -test are performed to verify the relationships between CBD and the influencing factors. The SMLR model has been used in many studies to link predictand and predictors to quantify the relative contributions of each variable (Tai et al. 2010; Zhang et al. 2014; Zhai et al. 2019; Pei et al. 2020). By using the SMLR, the standardized regression coefficients (Beta) can be compared among different variables, which represents the percentage of change in CBD with unique variations of each variable if all the other variables are unchanged after removing impacts of units and sample size (Wilks et al. 2006). In this study, the square of Beta is used as the explained variance; the closer this statistic is to 100%, the greater the contributions to the classified CBD.

The Earth System Modeling Framework is used for interpolating data into the $1^\circ \times 1^\circ$ grid to increase statistical robustness (Jones 1999). Besides, the K index is employed to represent the stratification instability of moist air in the lower and middle troposphere. The equation is as follows (Zhang et al. 2007):

$$K = (T_{850} - T_{500}) + T_{d850} - (T - T_d)_{700}.$$




Fig. 1 Location and altitudes of 378 meteorological stations in China

where T and T_d represent air temperature and dewpoint, and the subscript numbers show the geopotential height levels.

5 Results

5.1 Climatology and long-term trends of classified Chinese blue days in China

Figure 2a–c illustrate the spatial distribution of the average classified CBD in China during 1980–2018. At nationwide scale, the annual average CBD1–3 in China are 17.83, 82.01, and 62.00 days(d)/year(y), respectively. Basically, CBD1 exhibits a roughly geographic homogeneous pattern throughout the entire country, and the relatively low centers of CBD1 (below 5 days a year) are located in Yunnan, east of Tibet, and Hainan provinces. In contrast, Qinghai and most parts of Tibet share the maximum CBD1 with a value beyond 65 d/y. Compared to CBD1, the CBD2 tends to increase gradually from about 50d/y in southeast China to more than 80d/y in northwest China. As expected, almost all the low CBD2 centers reside in Southern China due to relatively higher low-cloud cover. The minimum CBD2 center, below 14 d/y, is located in Hebei province. CBD3 increases from southeast China to northwest China with the

maximum number in Xinjiang and eastern Inner Mongolia (over 70 d/y), and the minimum number in western Qinghai and southern Hebei (Fig. 2c, over 13.46 d/y, and 26.85 d/y).

The temporal evolutions of different CBD anomalies during 1980–2018 are shown in Fig. 2d. Upward trends are identified for CBD, CBD1, and CBD2 at a rate of 0.25, 0.34, and 0.39 days/year (d/y), respectively. Nevertheless, CBD3 is dominated by a declining trend exceeding -0.48 d/y, indicating a downward trend since 1980. Obvious increases of CBD1–2 can be found from 2014 to 2018, followed by an apparent decreasing CBD3 contemporaneous. This reveals the instant effect of emission controls on increasing CBD1–2 but an insufficient effect on increasing CBD3.

Figure 3 reveals the spatial distribution of trend coefficients with a horizontal resolution of $1^\circ \times 1^\circ$ in annual mean classified CBD from 1980 to 2018. It is obvious that CBD1 reveals an upward trend in almost 91.28% of grids (Fig. 3a). The most striking upward trend appears in northwestern China, with a value beyond 1 d/y. As for CBD2 (Fig. 3b), there is a similar dominant increasing trend over China (over 74.0%). Nearly all grids in the south of 38° N are dominated by an ascending trend, with a conspicuous center exceeding 0.95 d/y in southeastern China (SWC). In contrast, decreasing trends mostly occur in the north of 30° N, such as western Inner Mongolia, Qinghai, and some stations in northeast China. In contrast to CBD1 and CBD2, apart




Fig. 2 Spatial distribution of annual mean CBD under different grades in China during 1980–2018 (unit days), (a) CBD1, (b) CBD2, (c) CBD3, and (d) time series of the anomalous CBD in China from 1980 to 2018 (unit days)




Fig. 3 Observed distributions of trend coefficients (d/y) between 1980 and 2018 in China for (a) CBD1, (b) CBD2, (c) CBD3, and (d) CBD3 in winter; the white dots indicate the grid passing the 95% confidence level. The boxes overlaid indicate the six representative

regions, Northeast China (NEC), Northwest China (NWC), Yangtze-Huai River Valley (YHRV), Southeastern China (SWC), Pearl River Delta (PRD), and Beijing-Tianjin-Hebei (BTH)

from some regions in western China and northeast China, CBD3 displays a coherent significant decreasing trend (more than 82.7% of, Fig. 3c). A noteworthy downward trend can be found in eastern China, with the minimum value below -1.18 d/y.

Among the four seasons, the decrease of CBD3 is most prominent in winter. Fig. 3d shows the trend of wintertime CBD3 in China during 1980–2018, which is similar to the trend of annual CBD3. The regional mean winter CBD3 is decreasing at a rate of -0.11 d/y in China, with minimum centers located in northwestern China, southwestern China, and eastern China. These results are consistent with the increase in wintertime air pollution and an obvious increase in winter haze in China (Feng et al. 2020; Lu et al. 2020; Zhang et al. 2020b). In the following section, we focus on analyzing the reasons for the long-term change of wintertime CBD3 in China.

6 The reasons for wintertime CBD3 change in China

Four variables considered in our definition, including low cloud cover (LCC), visibility, relative humidity (RH), and non-precipitation day (non-REP) are taken into

consideration. The non-REP is calculated by counting the monthly number of non-rainy and non-snow days in winter. Table 1 shows the trend coefficients of selected meteorological variables and their explained variances (Beta square values). According to the CBD3 definition, there should be a positive relationship between visibility and CBD3, and a negative one between LCC and CBD3. The RH plays a more complicated role by affecting the optical characteristics of hydrophilic aerosols (Peng et al. 2009; Petters and Kreidenweis 2013).

In China, the growth of non-REP and visibility should contribute to the increase of CBD3. Nevertheless, reduced RH contributes to the decrease of the dry visibility (below -0.04 km/y) and finally reduces the CBD3 along with increasing LCC. The changes of RH, non-REP, LCC, and visibility account for 17.39%, 9.73%, 69.22%, and 1.04% of the area-mean CBD3 change in China, respectively.

Moreover, we divided China into different sub-regions, including northwest China (NWC), southwest China (SWC), the Yangtze-Huai River Valley (YHRV), Beijing-Tianjin-Hebei (BTH), Pearl River Delta (PRD), and northeast China (NEC). The locations of these sub-regions are shown in Fig. 2d. In NWC, with the most significant reduction below -0.45 d/y, the decrease of CBD3 is dominated by

Table 1 The trend coefficients of winter CBD3 and the associated meteorological parameters over China and some sub-regions, and the Beta square values are given in the right column (units: %)

Area	CBD3 (d/y)	RH (%/y)	Non-REP (d/y)	LCC (%/y)	Visibility (km/y)
China	−0.11	−0.06/17.39	0.18/9.73	0.09/69.22	0.018/1.042
YHRV	−0.42	−0.03/32.06	−0.02/0.67	0.05/69.89	−0.11/3.06
BTH	−0.02	−0.04/95.84	0.11/53.00	0.03/16.98	−0.04/3.42
PRD	0.02	−0.06/8.52	0.26/15.05	0.18/30.14	0.14/17.98
NEC	0.02	−0.07/24.30	0.20/30.58	0.15/0.03	0.14/1.99
SWC	−0.13	−0.11/14.67	0.32/22.00	−0.03/92.93	−0.08/0.90
NWC	−0.45	−0.00/5.95	0.06/0.76	0.20/46.24	−0.01/13.10

reducing visibility and increasing LCC concurrently, with explained variances over 13.10% and 46.24%, respectively. In BTH, under unfavorable RH, LCC, and visibility conditions, the growth of non-precipitation days leads to a decreasing trend below -0.02 d/y, with a dominant role of RH beyond 95.84%. In YHRV, with unfavorable situations for all variables, an obvious decreasing trend (over -0.42 d/y) is detected. The changes in RH and LCC play a more important role in this region, accounting for 32.06% and 69.89% of the variations of CBD3, respectively. In SWC, where there is a unique downward trend (below -0.03%) of LCC, the decrease of RH and visibility contribute to a reduced CBD3.

In other regions with a faint increase (over 0.02 d/y) like PRD and NEC, variations of RH, visibility, and non-REP contribute to deep blue days, with a leading role (more than 30.58%) of non-REP and LCC (over 30%) in NEC. It should be mentioned that considering the close relationships among variations of meteorological factors, the combined multiple explained variances may be greater than 100% due to multicollinearity. Therefore, these results only give reference information; the underlined mechanism should be analyzed by numerical model in the future.

To reveal the long-term variations of CBD3, we analyze meteorological and environmental factors for the change of visibility, RH, and LCC. Previous studies indicate that meteorological elements could have a crucial impact dynamically and thermodynamically (Zhang et al. 2014; Liu et al. 2017; Huang et al. 2020a; Ma et al. 2020). Therefore, air temperature at 2 m (t_{2m}), RH, WS, and K index are selected as major local meteorological conditions. The results are shown in Fig. 4a–d. A nationally downward trend of the K index in Qinghai, northwestern Sichuan, and south-eastern Tibet can be found in Fig. 4a. A more stable atmospheric stratification in the lower and middle troposphere favors formations of LCC in relatively moist near-surface conditions (Wu et al. 2013; Horton et al. 2014). Meanwhile, the decrease of the K index should be related to the increase of t_{2m} (Fig. 4b) and the decrease of RH (Fig. 4c). And it can partly explain the widespread increasing trend of CBD3, especially in coastal regions, consistent with previous studies (Cai et al. 2017; Li et al. 2018; Pei and Yan

2018). The wind speed also shows a significant decreasing trend in most parts of China (Fig. 4d). Previous studies show that decreasing wind speed can hinder the horizontal transportation of air pollutants (Zhang et al. 2014), which could partly explain the decrease of CBD3.

Earlier works find that through scattering and absorbing function, aerosol extinction can directly deteriorate visibility, and sulfate ammonium, organics, and nitrate ammonium are the largest contributors to the light extinction coefficient in China (Huang et al. 2012; Yang et al. 2012). Meanwhile, anthropogenic aerosols could affect clouds mainly as the most common condensation nuclei (CNN) in environments by altering the microphysical properties of liquid and ice clouds (Bierkens et al. 2001). Additionally, increasing CNN can inhibit low cloud precipitation by weakening the mixed-phase stratocumulus (Yun et al. 2009; Qian et al. 2010). As a result, there exists an intimate linkage between anthropogenic aerosols and CBD3. Then, the trend of main precursors of fine particles (including SO_2 , NO_x , and NMVOC) and BC (black carbon), which could affect visibility, are calculated. Only data from 1980 to 2014 are used due to data restriction.

Significant upward trends of SO_2 , NO_x , NMVOC, and BC are shown in Fig. 4e–h displays, with a rate of 0.21, 0.21, 0.12, and 0.01 Tg/y during 1980 and 2014, respectively. The major emission-increasing zones are accompanied by obvious decreasing trends of CBD3, with the correlation coefficients of -0.43 , -0.46 , -0.47 , and -0.51 , all passing the 99% confidence level. The result reveals a stronger connection between CBD3 and human activity than meteorology at a national scale, and the BC is the primary emission affecting CBD3 in China.

YHRV, which is the most populous region with the highest emission (at rates over 0.84, 0.84, 0.46, and 0.04 Tg/y for SO_2 , NO_x , NMVOC, and BC) in China, is found to be the most susceptible region. CBD3 in this region is significantly and negatively related to emissions, with the correlation coefficients all below -0.82 . In NWC, SWC, and BTH, similar situations can be found except for the major role of BC in NWC and SO_2 for another two, with a correlation coefficient of -0.39 , -0.25 , and -0.16 , respectively. The weak connections might indicate the complicated feedback




Fig. 4 The spatial distributions of trend coefficients of (a) K index, (b) t2m (°C/year), (c) relative humidity (%/year), (d) wind speed (m/s/year), (e) SO₂, (f) NO_x, (g) NMVOC, and (h) BC (units Tg/y) from

1980 to 2018 in winter. The white dots indicate the grid passing the 90% confidence level

between aerosols and meteorology, which needs further study. On the contrary, weak correlations in PRD and NEC might indicate a more important role of meteorological

conditions. We note that the results are probably affected by the limit of the analysis period. Further analysis is needed when the data are extended.

7 Conclusions and discussion

Although blue sky is a common weather phenomena and close to the needs of human, few studies have investigated it specially. In this study, we classify the types of blue sky according to meteorology condition (cloud cover and precipitation) and the concentrations of atmospheric aerosols (atmospheric visibility), and compute their long-term changes in China. It is found that deep blue sky days decrease significantly in China during 1980–2018, especially after 2013. NWC and YHRV share the most obvious downward trend on an annual scale or in winter.

Human activities at least partly contribute to the decrease of winter CBD3 in China. The variations of winter CBD3 are closely related to anthropogenic emissions of SO₂, NO_x, NMVOC, and BC with the correlation coefficients of −0.43, −0.46, −0.47, and −0.51, respectively. Meantime, the change of meteorological conditions is also responsible for the decrease of winter CBD3 in this period. The decrease in surface wind speed hinders air pollutant cleaning, and the increase in surface air temperature and decrease in relative humidity favor low cloud genesis, which is adverse to the occurrence of blue sky. These findings suggest that the occurrence of deep blue sky is affected by both the change of meteorological conditions and anthropogenic emissions.

This study only focus on China, but the definition of classified blue sky can extend to the global. Investigating how the blue sky change in the global may help us monitor the change of atmospheric environment of the earth, which deserve further study in the future.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00704-021-03898-1>.

Author contribution Gang Huang: conceptualization, supervision. Su Wang: methodology, software, writing-original draft, visualization, investigation. Kaiming Hu: writing-original draft, writing-reviewing, and editing, data curation. Lin Wang, Tie Dai, and Chunjiang Zhou: writing – reviewing.

Funding This work was supported by the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (2019QZKK0102), the National Natural Science Foundation of China (41831175, 91937302, 41721004, and 41775086), and the Key Deployment Project of the Centre for Ocean Mega-Research of Science, Chinese Academy of Sciences (COMS2019Q03).

Data availability The meteorological observations are archived by the China Meteorological Administration. The Re-Analysis data are provided by ECMWF. The air emissions species are provided by The Community Emissions Data System (CEDS).

Code availability The code used in this study will be available on reasonable request.

Declarations

Ethics approval The authors paid attention to the ethical rules in the study. There is no violation of ethics.

Consent to participate All the authors admitted that they have contributed to the study.

Consent for publication If this study is accepted, it can be published in the Theoretical and Applied Climatology journal.

Conflict of interest The authors declare no competing interests.

References

- Alexander LV, Zhang XB, Peterson TC, Caesar J, Vazquez-Aguirre JL (2006) Global observed changes in daily climate extremes of temperature and precipitation. *J Geophys Res-Atmos* 111:D05109
- Alexandersson H, Moberg A (1997) Homogenization of Swedish temperature data. Part I: homogeneity test for linear trends. *Int J Climatol* 17(1):25–34
- Bei N, Li X, Tie X, Zhao L, Wu J, Li X, Liu L, Shen Z, Li G (2020a) Impact of synoptic patterns and meteorological elements on the wintertime haze in the Beijing-Tianjin-Hebei region, China from 2013 to 2017. *Sci Total Environ* 704:135210
- Bei NF, Li XP, Tie XX, Zhao LN, Wu JR, Li X, Liu L, Shen ZX, Li GH (2020b) Impact of synoptic patterns and meteorological elements on the wintertime haze in the Beijing-Tianjin-Hebei region, China from 2013 to 2017. *Sci Total Environ* 704:12
- Bierkens MFP, Dolman H, Troch P (2001) Climate and the hydrological cycle
- Cai W, Li K, Liao H, Wang H, Wu L (2017) Weather conditions conducive to Beijing severe haze more frequent under climate?change. *Nat Clim Chang* 7(4):257–262
- Chang L, Wu Z, Xu J (2020) A comparison of haze pollution variability in China using haze indices based on observations. *Sci Total Environ* 715(May1):136929.1–136929.12
- China State Council (2013) Action plan on prevention and control of air pollution. http://www.gov.cn/zwggk/2013-09/12/content_2486773.htm. Accessed 12 Sep 2013
- China State Council (2018) the Three-year Action Plan for Blue-Sky Defense War. http://www.gov.cn/home/2018-07/03/content_5303211.htm. Accessed 03 Jul 2018
- Dai ZJ, Liu DY, Yu K, Cao L, Jiang YS (2020) Meteorological variables and synoptic patterns associated with air pollutions in Eastern China during 2013–2018. *Int J Environ Res Public Health* 17(7):19
- Dee D, Uppala SM, Simmons AJ (2011) The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *QJR Meteorol Soc* 137:553–597
- Ding AJ, Huang X, Nie W, Sun JN, Kerminen V-M, Petäjä T, Su H, Cheng YF, Yang X-Q, Wang MH (2016) Black carbon enhances haze pollution in megacities in China. *Geophys Res Lett* 43(6):2873–2879 n/a-n/a
- Dufour G, Eremenko M, Cuesta J, Doche C, Foret G, Beekmann M, Cheiney A, Wang Y, Cai Z, Liu Y, Takigawa M, Kanaya Y, Flaud JM (2015) Springtime daily variations in lower-tropospheric ozone over east Asia: the role of cyclonic activity and pollution as observed from space with IASI. *Atmos Chem Phys* 15(18):10839–10856
- Easterling DR, Meehl GA, Parmesan C, Changnon SA, Karl TR, Mearns LO (2019) Climate extremes: observations,

- modeling, and impacts. *Science* (New York, NY) 2000年 289卷 (期(5487)):2068–274页
- Fan GF, Ren L, Mao YD (2017) Reconstruction of haze day database based on the comparison between manual and automatic observations. *Zhongguo Huanjing Kexue/china Environ Sci* 37(4):1254–1261
- Fast JD, Gustafson WI, Easter RC, Zaveri RA, Peckham SE (2006) Evolution of Ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model. *J Geophys Res Atmos* 111(D21)
- Feng J, Liao H, Li YJ, Zhang ZY, Tang YX (2020) Long-term trends and variations in haze-related weather conditions in north China during 1980–2018 based on emission-weighted stagnation intensity. *Atmos Environ* 240:13
- Hoesly RM, Smith SJ, Feng LY, Klimont Z, Janssens-Maenhout G, Pitkanen T, Seibert JJ, Vu L, Andres RJ, Bolt RM, Bond TC, Dawidowski L, Kholod N, Kurokawa J, Li M, Liu L, Lu ZF, Moura MCP, O'Rourke PR, Zhang Q (2018) Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS). *Geosci Model Dev* 11(1):369–408
- Horton D, Skinner CB, Singh D, Diffenbaugh NS (2014) Occurrence and persistence of future atmospheric stagnation events. *Nat Clim Chang* 4(8):698
- Hu K, Huang G, Zheng XT, Xie SP, Qu X, Du Y, Liu L (2013) Interdecadal variations in ENSO influences on Northwest Pacific-East Asian early summertime climate simulated in CMIP5 models. *J Clim* 27(15):5982–5998
- Huang K, Zhuang G, Lin Y, Wang Q, Fu Q (2012) Impact of anthropogenic emission on air-quality over a megacity - revealed from an intensive atmospheric campaign during the Chinese Spring Festival. *Atmos Chem Phys* 12(7):17151–17185
- Huang RJ, He Y, Duan J, Li Y, O'Dowd CD (2020a) Contrasting sources and processes of particulate species in haze days with low and high relative humidity in wintertime Beijing. *Atmos Chem Phys* 20(14):9101–9114
- Huang X, Ding A, Gao J, Zheng B, Zhou D, Qi X, Tang R, Wang J, Ren C, Nie W, Chi X, Xu Z, Chen L, Li Y, Che F, Pang N, Wang H, Tong D, Qin W et al (2020b) Enhanced secondary pollution offset reduction of primary emissions during COVID-19 lockdown in China. *Natl Sci Rev* 8(2):nwaa137
- Huang X, Ding A, Wang Z, Ding K, Gao J, Chai F, Fu C (2020c) Amplified transboundary transport of haze by aerosol-boundary layer interaction in China. *Nat Geosci* 13(6):428–434
- Jones PW (1999) First- and second-order conservative remapping schemes for grids in spherical coordinates. *Mon Weather Rev* 127(9):2204–2210
- Kaiser DP (2002) Decreasing trends in sunshine duration over China for 1954–1998: Indication of increased haze pollution. *Geophysical Research Letters* 29(21)
- Kato N, Akimoto H (2007) Anthropogenic emissions of SO₂ and NO_x in Asia: Emission inventories. *Atmos Environ Part A* 26(16):2997–3017
- Li Z, Folmer H, Xue J (2014) To what extent does air pollution affect happiness? The case of the Jinchuan mining area, China. *Ecol Econ* 99(mar.):88–99
- Li QY, Zhang L, Wang T, Tham YJ, Ahmadov R, Xue LK, Zhang Q, Zheng JY (2016a) Impacts of heterogeneous uptake of dinitrogen pentoxide and chlorine activation on ozone and reactive nitrogen partitioning: improvement and application of the WRF-Chem model in southern China. *Atmos Chem Phys* 16(23):14875–14890
- Li Z, Lau W, Ramanathan V, Wu G, Ding Y, Manoj M, Liu J, Qian Y, Li J, Zhou T (2016b) Aerosol and monsoon climate interactions over Asia. *Rev Geophys* 54(4):866–929
- Li K, Liao H, Cai W, Yang Y (2018) Attribution of anthropogenic influence on atmospheric patterns conducive to recent most severe haze over Eastern China. *Geophys Res Lett* 45(4):2072–2081
- Lin M, Chan IN, Chan CY, Engling G, Bloss W (2012) Implications of regional surface ozone increases on visibility degradation in southeast China. *Tellus Series B-Chem Physical Meteorol* 64:19625
- Liu HR, Liu C, Xie ZQ, Li Y, Huang X, Wang SS, Xu J, Xie PH (2016) A paradox for air pollution controlling in China revealed by “APEC Blue” and “Parade Blue”. *Sci Rep* 6:1–13
- Liu HM, Fang CL, Zhang XL, Wang ZY, Bao C, Li FZ (2017) The effect of natural and anthropogenic factors on haze pollution in Chinese cities: a spatial econometrics approach. *J Clean Prod* 165:323–333
- Liu JY, Ren CH, Huang X, Nie W, Wang JP, Sun P, Chi XG, Ding AJ (2020) Increased aerosol extinction efficiency hinders visibility improvement in Eastern China. *Geophys Res Lett* 47(20):e2020GL090167
- Lu SH, Gong SL, He JJ (2020) Uncertainty analysis of spatiotemporal characteristics of haze pollution from 1961 to 2017 in China. *Atmos Pollut Res* 11(2):310–318
- Luehinger S (2010) Life satisfaction and transboundary air pollution. *Econ Lett* 107(1):4–6
- Ma N, Zhao CS, Chen J, Xu WY, Yan P, Zhou XJ (2014) A novel method for distinguishing fog and haze based on PM_{2.5}, visibility, and relative humidity. *Sci China-Earth Sci* 57(9):2156–2164
- Ma SM, Xiao ZM, Zhang YF, Wang LT, Shao M (2020) Assessment of meteorological impact and emergency plan for a heavy haze pollution episode in a core city of the North China Plain. *Aerosol Air Qual Res* 20(1):26–42
- Meinshausen M, Meinshausen N, Hare W, Raper SC, Frieler K, Knutti R, Frame DJ, Allen MR (2009) Greenhouse-gas emission targets for limiting global warming to 2 degrees C. *Nature* 458(7242):1158–1162
- Millar RJ, Fuglestad JS, Friedlingstein P, Grubb MJ, Allen MR (2017) Emission budgets and pathways consistent with limiting warming to 1.5°C. *Nat Geosci* 10(10):741–747
- Organization W M (2008) Guide of meteorological instruments and methods of observation. *Eos Trans* 55
- Panmao Z, Sun A, Ren F, Liu X, Gao B, Zhang Q (1999) Changes of climate extremes in China. *Clim Chang* 42:203–218
- Pei L, Yan Z (2018) Diminishing clear winter skies in Beijing towards a possible future. *Environ Res Lett* 13:124029
- Pei L, Yan Z, Sun Z, Miao S, Yao (2018) Increasing persistent haze in Beijing: potential impacts of weakening East Asian winter monsoons associated with northwestern Pacific sea surface temperature trends. *Atmos Chem Phys* 18:3173–3183
- Pei L, Yan Z, Chen D, Miao S (2020) Climate variability or anthropogenic emissions: which caused Beijing Haze? *Environ Res Lett* 15(3):034004
- Peng Y, Pan X, Jie T, Zhou X, Zhang R, Zeng LJP (2009) Hygroscopic growth of aerosol scattering coefficient: a comparative analysis between urban and suburban sites at winter in Beijing. *Particuology* 7(001):52–60
- Petters M, Kreidenweis SM (2013) A single parameter representation of hygroscopic growth and cloud condensation nucleus activity. *Atmos Chem Phys* 7(8):1081–1091
- Pope C, Burnett R, Thun M, Calle E, Krewski D, Ito K, Thurston G (2002) Lung Cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA : J Am Med Assoc* 287:1132–1141
- Qian DY, Gong D, Leung RJASL (2010) Light rain events change over North America, Europe, and Asia for 1973–2009. *Atmos Sci Lett* 11(4):301–306

- Sun YL, Wang ZF, Wild O, Xu WQ, Chen C, Fu PQ, Du W, Zhou LB, Zhang Q, Han TT, Wang QQ, Pan XL, Zheng HT, Li J, Guo XF, Liu JG, Worsnop DR (2016) "APEC Blue": secondary aerosol reductions from emission controls in Beijing. *Sci Rep* 6:1–9
- Tai A, Mickley LJ, Jacob DJ (2010) Correlations between fine particulate matter (PM_{2.5}) and meteorological variables in the United States: implications for the sensitivity of PM_{2.5} to climate change. *Atmos Environ* 44(32):3976–3984
- Wang S, Huang G, Lin J, Hu K, Gong H (2019) Chinese Blue days: a novel index and spatio-temporal variations. *Environ Res Lett* 14(7):074026
- Wang J, Liu Y, Ding Y (2020) On the connection between interannual variations of winter haze frequency over Beijing and different ENSO flavors. *Sci Total Environ* 740:140109
- Welsch H (2006) Environment and happiness: valuation of air pollution using life satisfaction data. *Ecol Econ* 58(4):801–813
- Wilks DS, Wilks DL, Wilks DS, Wilks DF, Wilks D, Wilks DL, Wilks D (2006) Statistical methods in the atmospheric science.
- Wu D (2004) A discuss on the difference between haze and fog and the warning of brownish haze weather. *Guangdong Meteorol* 4(1):1–4
- Wu P, Christidis N, Stott P (2013) Anthropogenic impact on Earth's hydrological cycle. *Nat Clim Chang* 3(9):807–810
- Xu ZJ, Shan W, Qi T, Gao J (2018) Characteristics of individual particles in Beijing before, during and after the 2014 APEC meeting. *Atmos Res* 203:254–260
- Yang Q, Wang YH, Zhao C, Liu Z, Gustafson WI, Shao M (2011) NO_x emission reduction and its effects on ozone during the 2008 Olympic games. *Environ Sci Technol* 45(15):6404–6410
- Yang L, Zhou X, Zhe W, Yang Z, Cheng S, Xu P, Gao X, Wei N, Wang X, Wang WJAE (2012) Airborne fine particulate pollution in Jinan, China: concentrations, chemical compositions and influence on visibility impairment. *Atmos Environ* 55(Aug.):506–514
- Yin Z, Wang H, Chen H (2017) Understanding severe winter haze events in the North China Plain in 2014: roles of climate anomalies. *Atmos Chem Phys* 17(3):1–27
- Yun Q, Gong D, Fan J, Leung LR, Bennartz R, Chen D, Wang W (2009) Heavy pollution suppresses light rain in China: Observations and modeling. *Journal of Geophysical Research: Atmospheres* 114
- Zhai SX, Jacob DJ, Wang X, Shen L, Li K, Zhang YZ, Gui K, Zhao TL, Liao H (2019) Fine particulate matter (PM_{2.5}) trends in China, 2013–2018: separating contributions from anthropogenic emissions and meteorology. *Atmos Chem Phys* 19(16):11031–11041
- Zhang C, Wei Z (2014) Panel estimation for income inequality and CO₂ emissions: a regional analysis in China. *Appl Energy* 136(C):382–392
- Zhang GC, Jiao MY, Li YX (2007) Techniques and methods of contemporary weather forecast (in Chinese). China Meteorological Press, Beijing, p 371
- Zhang RH, Li Q, Zhang RN (2014) Meteorological conditions for the persistent severe fog and haze event over eastern China in January 2013. *Sci China-Earth Sci* 57(1):26–35
- Zhang L, Wang T, Lv MY, Zhang Q (2015) On the severe haze in Beijing during January 2013: unraveling the effects of meteorological anomalies with WRF-Chem. *Atmos Environ* 104:11–21
- Zhang L, Shao JY, Lu X, Zhao YH, Hu YY, Henze DK, Liao H, Gong SL, Zhang Q (2016) Sources and Processes affecting fine particulate matter pollution over North China: an adjoint analysis of the Beijing APEC period. *Environ Sci Technol* 50(16):8731–8740
- Zhang L, Liu W, Hou K, Lin J, Song C, Zhou C, Huang B, Tong X, Wang J, Rhine W, Jiao Y, Wang Z, Ni R, Liu M, Zhang L, Wang Z, Yuebin W, Li X, Liu S, Wang Y (2019a) Air pollution exposure associates with increased risk of neonatal jaundice. *Nat Commun* 10:1–9
- Zhang L, Liu W, Hou K, Lin J, Zhou C, Tong X, Wang Z, Yuebin W, Jiang Y, Wang Z, Zheng Y, Lan Y, Liu S, Ni R, Liu M, Zhu P (2019b) Air pollution-induced missed abortion risk for pregnancies. *Nat Sustain* 2:1011–1017
- Zhang Q, Zheng Y, Tong D, Shao M, Hao JJ (2019c) Drivers of improved PM_{2.5} air quality in China from 2013 to 2017. *Proc Natl Acad Sci* 116(49):201907956
- Zhang Y, Gao L, Cao L, Yan Z, Wu Y (2020a) Decreasing atmospheric visibility associated with weakening winds from 1980 to 2017 over China. *Atmos Environ* 224:117314
- Zhang Y, Yin Z, Wang H (2020b) Roles of climate variability on the rapid increases of early winter haze pollution in North China after 2010. *Atmos Chem Phys* 20(20):12211–12221
- Zhou T, Gong D, Li J, Li B (2009) Detecting and understanding the multi-decadal variability of the East Asian Summer Monsoon – recent progress and state of affairs. *Meteorol Z* 18(4):455–467
- Zhu J, Liao H, Li J (2012) Increases in aerosol concentrations over eastern China due to the decadal-scale weakening of the East Asian summer monsoon. *Geophys Res Lett* 39(9):9809

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.